
**Monoclonal antibody** Catalog #: AMRe03775



# Summary

Phospho-GSK3 (Tyr216/Tyr279) Rabbit Monoclonal antibody **Production Name** 

Description Recombinant rabbit monoclonal antibody

Host Rabbit **Application** WB.IP Reactivity Mouse,Rat

### **Performance**

Conjugation Unconjugated Modification Phosphorylated

Isotype IgG

Clonality Monoclonal Antibody

**Form** Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% Buffer

protective protein

**Purification Affinity Purified** 

## **Immunogen**

**Gene Name** GSK3A/GSK3B

**Alternative Names** Serine/threonine-protein kinase GSK3A; Serine/threonine-protein kinase GSK3B

Gene ID 2931/2932

SwissProt ID P49840/P49841.

# **Application**

**Dilution Ratio** WB 1:500-1:1000,IP 1:20-1:50

**Molecular Weight** Calculated MW: 51 kDa; Observed MW: 47-51 kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

# Product Name: Phospho-GSK3 (Tyr216/Tyr279) Rabbit

Monoclonal antibody Catalog #: AMRe03775




# **Background**

Constitutively active protein kinase that acts as a negative regulator in the hormonal control of glucose homeostasis, Wnt signaling and regulation of transcription factors and microtubules, by phosphorylating and inactivating glycogen synthase (GYS1 or GYS2), CTNNB1/beta-catenin, APC and AXIN1. Requires primed phosphorylation of the majority of its substrates. Contributes to insulin regulation of glycogen synthesis by phosphorylating and inhibiting GYS1 activity and hence glycogen synthesis. Regulates glycogen metabolism in liver, but not in muscle. May also mediate the development of insulin resistance by regulating activation of transcription factors. In Wnt signaling, regulates the level and transcriptional activity of nuclear CTNNB1/beta-catenin. Facilitates amyloid precursor protein (APP) processing and the generation of APP-derived amyloid plaques found in Alzheimer disease. May be involved in the regulation of replication in pancreatic beta-cells. Is necessary for the establishment of neuronal polarity and axon outgrowth. Through phosphorylation of the anti-apoptotic protein MCL1, may control cell apoptosis in response to growth factors deprivation.

### Research Area

Neuroscience

# **Image Data**



Western blot analysis of Phospho-GSK3 (alpha/beta) (Tyr216/Tyr279) in C6, 3T3 lysates using Phospho-GSK3 (Tyr216/Tyr279) antibody.

# **Note**

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838