

# Product Name: Phospho-4E BP1 (Thr46) Rabbit Monoclonal Antibody Catalog #: AMRe04050

For research use only.

## **Summary**

**Description** Recombinant rabbit monoclonal antibody

Host Rabbit
Application WB,IHC,IP

Reactivity Human, Mouse, Rat
Conjugation Unconjugated
Modification Phosphorylated

**Isotype** IgG

Clonality Monoclonal
Form Liquid

Concentration 0.5mg/ml. The concentration of this product may be batch-dependent.

Storage Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

**Shipping** Ice bags

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% Buffer

protective protein

**Purification** Affinity Purification

# **Application**

**Dilution Ratio** WB 1:500-1:1000,IHC 1:50-1:100,IP 1:20-1:50

Molecular Weight Calculated MW: 13 kDa; Observed MW: 15-20 kDa

# **Antigen Information**

**Alternative Names** 

Gene Name EIF4EBP1

EIF4EBP1; Eukaryotic translation initiation factor 4E-binding protein 1; 4E-BP1; eIF4E-binding

protein 1; Phosphorylated heat- and acid-stable protein regulated by insulin 1; PHAS-I

**Gene ID** 1978

SwissProt ID Q13541

**Immunogen** A synthetic Phosphorylated peptide corresponding to residues target protein

# **Background**

Translation repressor protein 4E-BP1 (also known as PHAS-1) inhibits cap-dependent translation by binding to the translation

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




initiation factor eIF4E. Hyperphosphorylation of 4E-BP1 disrupts this interaction and results in activation of cap-dependent translation. Both the PI3 kinase/Akt pathway and FRAP/mTOR kinase regulate 4E-BP1 activity.


#### **Research Area**

**Epigenetics and Nuclear Signaling** 

## **Image Data**



Western blot analysis of Phospho-eIF4EBP1 (Thr46) in K562, C6, 3T3, Hela lysates using Phospho-4E BP1 (Thr46) antibody.



Immunohistochemistry analysis of paraffin-embedded Human breast cancer using eIF4EBP1/eIF4EBP2/eIF4EBP3 (Phospho- T45) antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838