
Product Name: Phospho-ErbB2(Y1221 + Y1222) (11Z13) Rabbit Monoclonal Antibody
Catalog #: AMRe05899

For research use only.

Summary

Description	Recombinant rabbit monoclonal antibody
Host	Rabbit
Application	WB
Reactivity	Human
Conjugation	Unconjugated
Modification	Phosphorylated
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Concentration	0.5mg/ml. The concentration of this product may be batch-dependent.
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:1000-1:5000
Molecular Weight	138kDa

Antigen Information

Gene Name	ERBB2 CD340; CerbB2; Erb b2 receptor tyrosine kinase 2; ERBB2; HER2; Herstatin; Human epidermal growth factor receptor 2; MLN19; NEU; NGL; Proto-oncogene Neu; Receptor tyrosine-protein kinase erbB-2; Tyrosine kinase type cell surface receptor HER2; V erb b2 avian erythroblastic leukemia viral oncogene homolog 2; V erb b2 avian erythroblastic leukemia viral oncoprotein 2;
Alternative Names	
Gene ID	2064.0
SwissProt ID	P04626

Immunogen

A synthetic phosphopeptide corresponding to residues surrounding Tyr1221/Tyr1222 of human ErbB 2


Background

Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization.

Research Area

Cancer

Image Data



Western blot analysis of Phospho-ErbB2(Y1221 + Y1222) expression in SKBR3 cell lysate.