Product Name: Phospho-S6K1 (T421 + S424) (1L11)

Rabbit Monoclonal Antibody

Catalog #: AMRe06002



Summary

Production Name Phospho-S6K1 (T421 + S424) (1L11) Rabbit Monoclonal Antibody

Description Rabbit Monoclonal Antibody

HostRabbitApplicationWB,IPReactivityHuman,Rat

Performance

ConjugationUnconjugatedModificationPhosphorylated

Isotype IgG

Clonality Monoclonal Form Liquid

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type

Buffer preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Purification Affinity purification

Immunogen

Gene Name RPS6KB1

EC 2.7.11.1, KS6B1, P70-S6K, RPS6KB1, Ribosomal protein S6 kinase, Ribosomal protein Alternative Names

S6 kinase, 70kDa, polypeptide 1, S6K, kinase p70S6K, p70-S6K

 Gene ID
 6198.0

 SwissProt ID
 P23443.

Application

Dilution Ratio WB 1:1000-1:2000,IP 1:20-1:50

Molecular Weight 59kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Phospho-S6K1 (T421 + S424) (1L11)

Rabbit Monoclonal Antibody

Catalog #: AMRe06002



Background

This gene encodes a member of the RSK (ribosomal S6 kinase) family of serine/threonine kinases. This kinase contains 2 non-identical kinase catalytic domains and phosphorylates several residues of the S6 ribosomal protein. Serine/threonineprotein kinase that acts downstream of mTOR signaling in response to growth factors and nutrients to promote cell proliferation, cell growth and cell cycle progression. Regulates protein synthesis through phosphorylation of EIF4B, RPS6 and EEF2K, and contributes to cell survival by repressing the pro-apoptotic function of BAD. Under conditions of nutrient depletion, the inactive form associates with the EIF3 translation initiation complex. Upon mitogenic stimulation, phosphorylation by the mammalian target of rapamycin complex 1 (mTORC1) leads to dissociation from the EIF3 complex and activation. The active form then phosphorylates and activates several substrates in the pre-initiation complex, including the EIF2B complex and the cap-binding complex component EIF4B. Also controls translation initiation by phosphorylating a negative regulator of EIF4A, PDCD4, targeting it for ubiquitination and subsequent proteolysis. Promotes initiation of the pioneer round of protein synthesis by phosphorylating POLDIP3/SKAR. In response to IGF1, activates translation elongation by phosphorylating EEF2 kinase (EEF2K), which leads to its inhibition and thus activation of EEF2. Also plays a role in feedback regulation of mTORC2 by mTORC1 by phosphorylating RICTOR, resulting in the inhibition of mTORC2 and AKT1 signaling. Mediates cell survival by phosphorylating the pro-apoptotic protein BAD and suppressing its pro-apoptotic function. Phosphorylates mitochondrial URI1 leading to dissociation of a URI1-PPP1CC complex. The free mitochondrial PPP1CC can then dephosphorylate RPS6KB1 at Thr-412, which is proposed to be a negative feedback mechanism for the RPS6KB1 anti-apoptotic function. Mediates TNF-alpha-induced insulin resistance by phosphorylating IRS1 at multiple serine residues, resulting in accelerated degradation of IRS1. In cells lacking functional TSC1-2 complex, constitutively phosphorylates and inhibits GSK3B. May be involved in cytoskeletal rearrangement through binding to neurabin. Phosphorylates and activates the pyrimidine biosynthesis enzyme CAD, downstream of MTOR (PubMed:11500364, PubMed:12801526, PubMed:14673156, PubMed:15071500, PubMed: 15341740, PubMed:16286006, PubMed:17052453, PubMed:17053147, PubMed:17936702, PubMed:18952604, PubMed:19085255, PubMed:19720745, PubMed: 19935711, PubMed:<a

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Phospho-S6K1 (T421 + S424) (1L11)

Rabbit Monoclonal Antibody


Catalog #: AMRe06002



href="http://www.uniprot.org/citations/19995915" target="_blank">19995915, PubMed:23429703). Following activation by mTORC1, phosphorylates EPRS and thereby plays a key role in fatty acid uptake by adipocytes and also most probably in interferongamma-induced translation inhibition (PubMed:28178239).

Research Area

Image Data



Western blot analysis of SK61 phosphorylation expression in HEK293 cell lysate.

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838