

## **Product Name: CD10 (15I14) Rabbit Monoclonal Antibody**

Catalog #: AMRe08178

For research use only.

#### **Summary**

**Description** Recombinant rabbit monoclonal antibody

**Host** Rabbit

Application WB,IHC,ICC/IF,IP

Reactivity Human,Mouse,Rat

Conjugation Unconjugated

Modification Unmodified

**Isotype** IgG

Clonality Monoclonal
Form Liquid

**Concentration** 0.3mg/ml. The concentration of this product may be batch-dependent. **Storage** Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

**Shipping** Ice bags

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% New type preservative

Buffer N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw

cycle.

**Purification** Affinity purification

### **Application**

**Dilution Ratio** WB 1:500-1:2000,IHC 1:200-1:1000,ICC/IF 1:20-1:50,IP 1:20-1:50

Molecular Weight 86kDa

### **Antigen Information**

Gene Name MME

Atriopeptidase; CALLA; CD10; CD10 antigen; Enkephalinase; EPN; Membrane metallo

Alternative Names endopeptidase (neutral endopeptidase, enkephalinase, CALLA, CD10); CALLA CD10; NEP;

Neprilysin; Neutral endopeptidase;

 Gene ID
 4311.0

 SwissProt ID
 P08473

**Immunogen** A synthetic peptide of human CD10

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




# **Background**

CD10 is a transmembrane type II molecule and functions as a metallo-peptidase requiring zinc. Specifically, CD10 cleaves 1-3 amino-terminal amino acids from peptides with a preference for neutral amino acids (valine, iso-leucine, phenylalanine, leucine or alanine). Involved in the degradation of atrial natriuretic factor (ANF). Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30 amino acids (PubMed:15283675, PubMed:8168535). Biologically important in the destruction of opioid peptides such as Met- and Leu-enkephalins by cleavage of a Gly-Phe bond (PubMed:17101991). Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9 (PubMed:15283675). Involved in the degradation of atrial natriuretic factor (ANF) and brain natriuretic factor (BNP(1- 32)) (PubMed:2531377, PubMed:2972276, PubMed:16254193). Displays UV-inducible elastase activity toward skin preelastic and elastic fibers (PubMed:20876573).

#### **Research Area**

Immunology

#### **Image Data**



Western blot analysis of CD10 expression in Ramos cell lysate.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838