

Product Name: CD163 (103) Rabbit Monoclonal Antibody**Catalog #: AMRe08240**

For research use only.

Summary

Description	Recombinant rabbit monoclonal antibody
Host	Rabbit
Application	WB,IHC,IF-P
Reactivity	Human
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Concentration	0.3mg/ml. The concentration of this product may be batch-dependent.
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:200-1:2000,IF-P 1:200-1:2000
Molecular Weight	125kDa

Antigen Information

Gene Name	CD163
Alternative Names	CD163; CD163 antigen; CD163 molecule; SCAR11; sCD163; Soluble CD163;
Gene ID	9332.0
SwissProt ID	Q86VB7
Immunogen	Recombinant protein of human CD163

Background


Involved in clearance and endocytosis of hemoglobin/haptoglobin complexes by macrophages and may thereby protect

tissues from free hemoglobin-mediated oxidative damage. May play a role in the uptake and recycling of iron, via endocytosis of hemoglobin/haptoglobin and subsequent breakdown of heme. Binds hemoglobin/haptoglobin complexes in a calcium-dependent and pH-dependent manner. Acute phase-regulated receptor involved in clearance and endocytosis of hemoglobin/haptoglobin complexes by macrophages and may thereby protect tissues from free hemoglobin-mediated oxidative damage. May play a role in the uptake and recycling of iron, via endocytosis of hemoglobin/haptoglobin and subsequent breakdown of heme. Binds hemoglobin/haptoglobin complexes in a calcium-dependent and pH-dependent manner. Exhibits a higher affinity for complexes of hemoglobin and multimeric haptoglobin of HP*1F phenotype than for complexes of hemoglobin and dimeric haptoglobin of HP*1S phenotype. Induces a cascade of intracellular signals that involves tyrosine kinase-dependent calcium mobilization, inositol triphosphate production and secretion of IL6 and CSF1. Isoform 3 exhibits the higher capacity for ligand endocytosis and the more pronounced surface expression when expressed in cells.

Research Area

Immunology; Innate Immunity; Macrophage / Inflamm; Cell Type Markers; CD; Myeloid Cells; Cardiovascular; Atherosclerosis; Vascular Inflammation; Innate and adaptive immunity

Image Data



Western blot analysis of CD163 expression in Human fetal kidney lysate.