
Product Name: E2F2 (6C14) Rabbit Monoclonal Antibody**Catalog #: AMRe10252**

For research use only.

Summary

Description	Recombinant rabbit monoclonal antibody
Host	Rabbit
Application	WB,FC
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Concentration	0.5mg/ml. The concentration of this product may be batch-dependent.
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:1000-1:5000,FC 1:100-1:500
Molecular Weight	48kDa

Antigen Information

Gene Name	E2F2
Alternative Names	De2f2; E2F transcription factor 2; E2F-2; Transcription factor E2F2;
Gene ID	1870.0
SwissProt ID	Q14209
Immunogen	A synthetic peptide of human E2F2


Background

Play an important role in the negative regulation of cell proliferation. Functional inactivation of Rb can be mediated either

through mutation or as a consequence of interaction with DNA tumor virus-encoded proteins. The DRTF1/E2F complex functions in the control of cell-cycle progression from g1 to s phase. E2F2 binds specifically to RB1 in a cell-cycle dependent manner. Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The DRTF1/E2F complex functions in the control of cell-cycle progression from g1 to s phase. E2F2 binds specifically to RB1 in a cell-cycle dependent manner.

Research Area

Image Data



Western blot analysis of E2F2 expression in K562 cell lysate.