
Product Name: Furin (6Z15) Rabbit Monoclonal Antibody**Catalog #: AMRe11184**

For research use only.

Summary

Description	Recombinant rabbit monoclonal antibody
Host	Rabbit
Application	WB,IHC,ICC/IF
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Concentration	0.5mg/ml. The concentration of this product may be batch-dependent.
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:1000-1:5000,IHC 1:50-1:100,ICC/IF 1:100-1:200
Molecular Weight	87kDa

Antigen Information

Gene Name	FURIN
Alternative Names	FUR; FURIN; PACE; PCSK3; SPC1;
Gene ID	5045.0
SwissProt ID	P09958
Immunogen	Recombinant protein of human Furin


Background

Furin is likely to represent the ubiquitous endoprotease activity within constitutive secretory pathways and capable of cleavage

at the RX(K/R)R consensus motif. Ubiquitous endoprotease within constitutive secretory pathways capable of cleavage at the RX(K/R)R consensus motif (PubMed:11799113, PubMed:1629222, PubMed:1713771, PubMed:2251280, PubMed:24666235, PubMed:25974265, PubMed:7592877, PubMed:7690548, PubMed:9130696). Mediates processing of TGFB1, an essential step in TGF-beta-1 activation (PubMed:7737999). Converts through proteolytic cleavage the non-functional Brain natriuretic factor prohormone into its active hormone BNP(1-32) (PubMed:20489134, PubMed:21763278).

Research Area

Image Data



Western blot analysis of Furin expression in HepG2 cell lysate.