Product Name: HTRA2 (8W8) Rabbit Monoclonal

Antibody

Catalog #: AMRe12276



Summary

Production Name HTRA2 (8W8) Rabbit Monoclonal Antibody

Description Rabbit Monoclonal Antibody

Host Rabbit

Application WB,IHC-P,IP,IF-P **Reactivity** Human,Mouse,Rat

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

Clonality Monoclonal Form Liquid

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type

Buffer preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Purification Affinity purification

Immunogen

Gene Name HTRA2

HTRA2; HtrA-like serine protease; OMI; PARK13; Protease; PRSS25; Serine protease 25; Alternative Names

HtrA serine peptidase 2; Serine proteinase OMI;

Gene ID 27429.0 **SwissProt ID** 043464.

Application

Dilution Ratio WB 1:1000-1:2000, IHC-P/IF-P 1:100, IP 1:20

Molecular Weight 49kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: HTRA2 (8W8) Rabbit Monoclonal

Antibody

Catalog #: AMRe12276




Background

High temperature requirement protein A2 (HtrA2)/Omi is a serine protease with homology to the E. coli HtrA protein (DegP) and is thought to be involved in apoptosis and stress-induced degradation of misfolded proteins. While HtrA2 was originally identified to be present in either the nucleus or endoplasmic reticulum, subsequent studies have shown that it localizes in mitochondria and is released during apoptosis. Serine protease that shows proteolytic activity against a non-specific substrate beta-casein. Promotes or induces cell death either by direct binding to and inhibition of BIRC proteins (also called inhibitor of apoptosis proteins, IAPs), leading to an increase in caspase activity, or by a BIRC inhibition-independent, caspase- independent and serine protease activity-dependent mechanism. Cleaves THAP5 and promotes its degradation during apoptosis. Isoform 2 seems to be proteolytically inactive.

Research Area

Image Data



Western blot analysis of HTRA2 expression in Jurkat cell lysate.

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838