

---

**Product Name: Interferon Receptor alpha (15I18) Rabbit Monoclonal Antibody****Catalog #: AMRe12685**

For research use only.

**Summary**

<b>Description</b>	Recombinant rabbit monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB
<b>Reactivity</b>	Human
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Concentration</b>	0.5mg/ml. The concentration of this product may be batch-dependent.
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Purification</b>	Affinity purification

**Application**

<b>Dilution Ratio</b>	WB 1:1000-1:5000
<b>Molecular Weight</b>	64kDa

**Antigen Information**

<b>Gene Name</b>	IFNAR1
<b>Alternative Names</b>	AVP; IFN alpha REC; IFNAR1; IFNBR; IFRC; interferon (alpha beta and omega) receptor 1;
<b>Gene ID</b>	3454.0
<b>SwissProt ID</b>	P17181
<b>Immunogen</b>	A synthetic peptide of human Interferon alpha/beta receptor 1

**Background**


Interferon Receptor alpha (IFN- $\alpha$  R1) is a class II cytokine receptor which belongs to type I human interferons (IFNs) family. IFNs

plays a role in antiviral, antiproliferative, immunomodulatory, antitumor, and antiparasitic activities by inducing transcription of IFN-stimulated genes (ISGs) through activation of the Jak-STAT pathway. Component of the receptor for type I interferons, including interferons alpha, IFNB1 and IFNW1 (PubMed:2153461, PubMed:7665574, PubMed:10049744, PubMed:14532120, PubMed:15337770, PubMed:21854986). Functions in general as heterodimer with IFNAR2 (PubMed:7665574, PubMed:10049744, PubMed:21854986). Type I interferon binding activates the JAK-STAT signaling cascade, and triggers tyrosine phosphorylation of a number of proteins including JAKs, TYK2, STAT proteins and the IFNR alpha- and beta-subunits themselves (PubMed:7665574, PubMed:21854986, PubMed:32972995). Can form an active IFNB1 receptor by itself and activate a signaling cascade that does not involve activation of the JAK-STAT pathway (By similarity).

## Research Area

Immunology

## Image Data



Western blot analysis of Interferon Receptor alpha expression in SH-SY5Y cell lysate.