
Product Name: KAT8 / MYST1 / MOF (9E16) Rabbit Monoclonal Antibody**Catalog #: AMRe12907**

For research use only.

Summary

Description	Recombinant rabbit monoclonal antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,FC,IP
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Concentration	0.5mg/ml. The concentration of this product may be batch-dependent.
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:200-1:500,ICC/IF 1:500-1:2000,FC 1:100-1:200,IP 1:100-1:200
Molecular Weight	52kDa

Antigen Information

Gene Name	KAT8
Alternative Names	hMOF; MOF; MOZ; MYST1; SAS2; SAS3; YBF2; YBF2/SAS3; ZC2HC8;
Gene ID	84148.0
SwissProt ID	Q9H7Z6
Immunogen	A synthetic peptide of human MOF


Background

Histone acetyltransferase which may be involved in transcriptional activation. May influence the function of ATM. Histone

acetyltransferase which may be involved in transcriptional activation (PubMed:12397079, PubMed:22020126). May influence the function of ATM (PubMed:15923642). As part of the MSL complex it is involved in acetylation of nucleosomal histone H4 producing specifically H4K16ac (PubMed:16227571, PubMed:16543150, PubMed:21217699, PubMed:22547026, PubMed:22020126). As part of the NSL complex it may be involved in acetylation of nucleosomal histone H4 on several lysine residues (PubMed:20018852, PubMed:22547026). That activity is less specific than the one of the MSL complex (PubMed:20018852, PubMed:22547026). Can also acetylate TP53/p53 at 'Lys-120'.

Research Area

Image Data



Western blot analysis of KAT8 / MYST1 / MOF expression in HeLa cell lysate.