

Product Name: RAB8A (15Q17) Rabbit Monoclonal Antibody
Catalog #: AMRe16803

Summary

Production Name	RAB8A (15Q17) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,IP
Reactivity	Human

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	RAB8A
Alternative Names	MEL; Mel transforming oncogene; Oncogene c mel; RAB8; RAB8A;
Gene ID	4218.0
SwissProt ID	P61006.

Application

Dilution Ratio	WB 1:1000, IP 1:20
Molecular Weight	24kDa

Product Name: RAB8A (15Q17) Rabbit Monoclonal Antibody
Catalog #: AMRe16803




Background

May be involved in vesicular trafficking and neurotransmitter release. Together with RAB11A, RAB3IP, the exocyst complex, PARD3, PRKCI, ANXA2, CDC42 and DNMBP promotes transcytosis of PODXL to the apical membrane initiation sites (AMIS), apical surface formation and lumenogenesis. Together with MYO5B and RAB11A participates in epithelial cell polarization. The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. That Rab is involved in polarized vesicular trafficking and neurotransmitter release. Together with RAB11A, RAB3IP, the exocyst complex, PARD3, PRKCI, ANXA2, CDC42 and DNMBP promotes transcytosis of PODXL to the apical membrane initiation sites (AMIS), apical surface formation and lumenogenesis (PubMed: [20890297](http://www.uniprot.org/citations/20890297)). Together with MYO5B and RAB11A participates in epithelial cell polarization (PubMed: [21282656](http://www.uniprot.org/citations/21282656)). May be involved in ciliogenesis (PubMed: [21844891](http://www.uniprot.org/citations/21844891), PubMed: [30398148](http://www.uniprot.org/citations/30398148)). Together with MICALL2, may also regulate adherens junction assembly (By similarity). May play a role in insulin-induced transport to the plasma membrane of the glucose transporter GLUT4 and therefore play a role in glucose homeostasis (By similarity). Involved in autophagy (PubMed: [27103069](http://www.uniprot.org/citations/27103069)).

Research Area

Image Data



Western blot analysis of RAB8A expression in HeLa cell lysate.

Note

**Product Name: RAB8A (15Q17) Rabbit Monoclonal
Antibody**
Catalog #: AMRe16803



For research use only.