

---

**Product Name: RPE65 (5T1) Rabbit Monoclonal Antibody****Catalog #: AMRe17356**

For research use only.

**Summary**

<b>Description</b>	Recombinant rabbit monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC,IP,IF-P
<b>Reactivity</b>	Human,Mouse,Rat
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Concentration</b>	0.25mg/ml. The concentration of this product may be batch-dependent.
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Purification</b>	Affinity purification

**Application**

<b>Dilution Ratio</b>	WB 1:500-1:2000,IHC 1:500-1:2000,IP 1:20-1:50,IF-P 1:500-1:2000
<b>Molecular Weight</b>	61kDa

**Antigen Information**


<b>Gene Name</b>	RPE65
<b>Alternative Names</b>	LCA2; mRPE65; p63; rd12; Retinal pigment epithelium specific 61 kDa protein; RP20; RPE65; sRPE65;
<b>Gene ID</b>	6121.0
<b>SwissProt ID</b>	Q16518
<b>Immunogen</b>	A synthetic peptide of human RPE65

**Background**

Plays important roles in the production of 11-cis retinal and in visual pigment regeneration. The soluble form binds vitamin A (all-trans-retinol), making it available for LRAT processing to all-trans-retinyl ester. The membrane form, palmitoylated by LRAT, binds all-trans-retinyl esters, making them available for IMH (isomerohydrolase) processing to all-cis-retinol. Critical isomerohydrolase in the retinoid cycle involved in regeneration of 11-cis-retinal, the chromophore of rod and cone opsins. Catalyzes the cleavage and isomerization of all-trans-retinyl fatty acid esters to 11-cis-retinol which is further oxidized by 11-cis retinol dehydrogenase to 11-cis-retinal for use as visual chromophore (PubMed:16116091). Essential for the production of 11-cis retinal for both rod and cone photoreceptors (PubMed:17848510). Also capable of catalyzing the isomerization of lutein to meso-zeaxanthin an eye- specific carotenoid (PubMed:28874556). The soluble form binds vitamin A (all-trans-retinol), making it available for LRAT processing to all- trans-retinyl ester. The membrane form, palmitoylated by LRAT, binds all-trans-retinyl esters, making them available for IMH (isomerohydrolase) processing to all-cis-retinol. The soluble form is regenerated by transferring its palmitoyl groups onto 11-cis-retinol, a reaction catalyzed by LRAT (By similarity).

## Research Area

### Image Data



Western blot analysis of RPE65 expression in mouse eyeball lysate.