

Product Name: S100P (5R5) Rabbit Monoclonal Antibody**Catalog #: AMRe17476**

For research use only.

Summary

Description	Recombinant rabbit monoclonal antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,FC,IP
Reactivity	Human
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Concentration	0.33mg/ml. The concentration of this product may be batch-dependent.
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:1000-1:5000,IHC 1:200-1:2000,ICC/IF 1:200-1:500,FC 1:200-1:1000,IP 1:20-1:50
Molecular Weight	10kDa

Antigen Information

Gene Name	S100P
Alternative Names	Protein S100-E; Protein S100P; S100 calcium binding protein P; S100E;
Gene ID	6286.0
SwissProt ID	P25815
Immunogen	A synthetic peptide of human S100P


Background

S100 proteins regulate a variety of cellular processes such as cell growth and motility, cell cycle progression, transcription, and

differentiation. S100P, a member of the S100 family, is a 95 amino acid protein. Its expression is increased in a number of tumors including pancreas, lung, breast, and ovary carcinomas in response to transcriptional activation. May function as calcium sensor and contribute to cellular calcium signaling. In a calcium-dependent manner, functions by interacting with other proteins, such as EZR and PPP5C, and indirectly plays a role in physiological processes like the formation of microvilli in epithelial cells. May stimulate cell proliferation in an autocrine manner via activation of the receptor for activated glycation end products (RAGE).

Research Area

Image Data



Western blot analysis of S100P expression in SW480 cell lysate.