

Product Name: Torsin A (2F5) Rabbit Monoclonal Antibody**Catalog #: AMRe19137**

For research use only.

Summary

Description	Recombinant rabbit monoclonal antibody
Host	Rabbit
Application	WB,FC
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Concentration	0.5mg/ml. The concentration of this product may be batch-dependent.
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,FC 1:50-1:200
Molecular Weight	38kDa

Antigen Information

Gene Name	TOR1A
Alternative Names	TOR1A;DQ2; DYT1; TorsinA;
Gene ID	1861.0
SwissProt ID	O14656
Immunogen	A synthetic peptide of human Torsin A/DYT1


Background

The neurological condition Dystonia is associated with sustained muscle contractions and abnormal posturing. TorsinA,

torsinB, torp2A and torp3A belong to the family of ATPases associated with cellular activities (AAA+) and mutations in torsinA cause early onset dystonia. TorsinA has been shown to suppress intracellular protein aggregation in *C. elegans* and possesses chaperon activity. Interestingly, torsinA is highly expressed in dopaminergic neurons and associates with alpha-synuclein in Lewy bodies, which pathologically characterize Parkinson's Disease. Protein with chaperone functions important for the control of protein folding, processing, stability and localization as well as for the reduction of misfolded protein aggregates. Involved in the regulation of synaptic vesicle recycling, controls STON2 protein stability in collaboration with the COP9 signalosome complex (CSN). In the nucleus, may link the cytoskeleton with the nuclear envelope, this mechanism seems to be crucial for the control of nuclear polarity, cell movement and, specifically in neurons, nuclear envelope integrity. Participates in the cellular trafficking and may regulate the subcellular location of multipass membrane proteins such as the dopamine transporter SLC6A3, leading to the modulation of dopamine neurotransmission. In the endoplasmic reticulum, plays a role in the quality control of protein folding by increasing clearance of misfolded proteins such as SGCE variants or holding them in an intermediate state for proper refolding. May have a redundant function with TOR1B in non- neural tissues.

Research Area

Image Data



Western blot analysis of Torsin A expression in 293T cell lysate.