

Product Name: Sumo 1 Rabbit Monoclonal Antibody

Catalog #: AMRe21347

For research use only.

Summary

Description	Recombinant rabbit monoclonal antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA,IP
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG,Kappa
Clonality	Monoclonal
Form	Liquid
Concentration	0.3mg/ml. The concentration of this product may be batch-dependent.
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%protective protein
Purification	Protein A

Application

Dilution Ratio	WB 1:2000-1:10000,IHC 1:2000-1:10000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200
Molecular Weight	Calculated MW:12kD;Observed MW:80kD

Antigen Information


Gene Name	SUMO1 SUMO1;SMT3C;SMT3H3;UBL1;OK/SW-cl.43;Small ubiquitin-related modifier 1;SUMO-
Alternative Names	1;GAP-modifying protein 1;GMP1;SMT3 homolog 3;Sentrin;Ubiquitin-homology domain protein PIC1;Ubiquitin-like protein SMT3C;Smt3C;Ubiquitin-like protein
Gene ID	7341
SwissProt ID	P63165
Immunogen	A synthetic peptide of human Sumo 1

Background


Cell localization: Cytoplasm, Nucleus. This gene encodes a protein that is a member of the SUMO (small ubiquitin-like modifier) protein family. It functions in a manner similar to ubiquitin in that it is bound to target proteins as part of a post-translational modification system. However, unlike ubiquitin which targets proteins for degradation, this protein is involved in a variety of cellular processes, such as nuclear transport, transcriptional regulation, apoptosis, and protein stability. It is not active until the last four amino acids of the carboxy-terminus have been cleaved off. Several pseudogenes have been reported for this gene. Alternate transcriptional splice variants encoding different isoforms have been characterized. [provided by RefSeq, Jul 2008],

Research Area

Image Data



Western blot analysis of lysates from Jurkat cells, using Sumo 1 Rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG antibody was used to detect the antibody.



Immunohistochemical analysis of paraffin-embedded Human lung carcinoma tissue. 1, Sumo 1 Rabbit Monoclonal Antibody was diluted at 1:200 (4°C, overnight). 2, EDTA pH 9.0 was used for antibody retrieval (>98°C, 20min). 3, Secondary antibody was diluted at 1:200 (room temperature, 30min).