# **Product Name: Cdc2 Rabbit Monoclonal Antibody**

Catalog #: AMRe21356



### **Summary**

Production Name Cdc2 Rabbit Monoclonal Antibody

**Description** Rabbit Monoclonal Antibody

**Host** Rabbit

**Application** WB,IHC,IF,IP,ELISA **Reactivity** Human,Mouse

#### **Performance**

ConjugationUnconjugatedModificationUnmodifiedIsotypeIgG,KappaClonalityMonoclonalFormLiquid

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

**Buffer** PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protective protein

**Purification** Protein A

## **Immunogen**

Gene Name CDK1

CDK1;CDC2;CDC28A;CDKN1;P34CDC2;Cyclin-dependent kinase 1;CDK1;Cell division Alternative Names

control protein 2 homolog; Cell division protein kinase 1;p34 protein kinase

Gene ID 983

SwissProt ID P06493.

# **Application**

IHC 1:1000-1:4000;WB 1:2000-1:10000;IF 1:200-1:1000;ELISA 1:5000-1:20000;IP 1:50-

**Dilution Ratio** 

1:200;

Molecular Weight Calculated MW:34kD;Observed MW:34kD

# **Background**


# Product Name: Cdc2 Rabbit Monoclonal Antibody Catalog #: AMRe21356



Cell localization:Cytoplasm, Nucleus.cyclin dependent kinase 1(CDK1) Homo sapiens The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2009],

#### **Research Area**

#### **Image Data**



Western blot analysis of lysates from K562

cells, using Cdc2 Rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG antibody was used to detect the antibody.

#### Note

For research use only.