

Product Name: eIF2 α Rabbit Monoclonal Antibody**Catalog #: AMRe21372**

For research use only.

Summary

Description	Recombinant rabbit monoclonal antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA,IP
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG,Kappa
Clonality	Monoclonal
Form	Liquid
Concentration	0.3mg/ml. The concentration of this product may be batch-dependent.
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%protective protein
Purification	Protein A

Application

Dilution Ratio	WB 1:2000-1:10000,IHC 1:1000-1:5000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200
Molecular Weight	Calculated MW:36kD;Observed MW:36kD

Antigen Information

Gene Name	EIF2S1
Alternative Names	EIF2S1;EIF2A;Eukaryotic translation initiation factor 2 subunit 1;Eukaryotic translation initiation factor 2 subunit alpha;eIF-2-alpha;eIF-2A;eIF-2alpha
Gene ID	1965
SwissProt ID	P05198
Immunogen	A synthetic peptide corresponding to target protein


Background

Cell localization:Cytoplasm.The translation initiation factor EIF2 catalyzes the first regulated step of protein synthesis initiation,

promoting the binding of the initiator tRNA to 40S ribosomal subunits. Binding occurs as a ternary complex of methionyl-tRNA, EIF2, and GTP. EIF2 is composed of 3 nonidentical subunits, the 36-kD EIF2-alpha subunit (EIF2S1), the 38-kD EIF2-beta subunit (EIF2S2; MIM 603908), and the 52-kD EIF2-gamma subunit (EIF2S3; MIM 300161). The rate of formation of the ternary complex is modulated by the phosphorylation state of EIF2-alpha (Ernst et al., 1987 [PubMed 2948954]).[supplied by OMIM, Feb 2010],

Research Area

Image Data



Western blot analysis of lysates from MCF7 cells, using eIF2α Rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG antibody was used to detect the antibody.