
Product Name: CD44 Rabbit Monoclonal Antibody**Catalog #: AMRe21424**

For research use only.

Summary

Description	Recombinant rabbit monoclonal antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA,IP
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG,Kappa
Clonality	Monoclonal
Form	Liquid
Concentration	0.3mg/ml. The concentration of this product may be batch-dependent.
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%protective protein
Purification	Protein A

Application

Dilution Ratio	WB 1:1000-1:5000,IHC 1:20000-1:50000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200
Molecular Weight	Calculated MW:81kD;Observed MW:81kD

Antigen Information

Gene Name	CD44
Alternative Names	LHR antibody;BA-1 antibody;CD 44 antibody;CD44 antibody;CD44 antigen antibody;CD44 molecule;Indian blood group;antibody;CD44 molecule antibody;CD44_HUMAN antibody;CDw44 antibody;CDW44 antigen antibody;Cell surface glycoprotein CD44 antibody;chondroitin sulfate proteoglycan 8 antibody;CSPG8 antibody;ECMR-III antibody;Epican antibody;Extracellular matrix receptor III antibody;GP90 lymphocyte homing/adhesion receptor antibody;HCELL antibody;hematopoietic cell E- and L-selectin ligand antibody;Heparan sulfate proteoglycan antibody;Hermes antigen antibody;homing function and Indian blood group system antibody;HSA antibody;HUTCH-I antibody;HUTCH1

antibody;HUTCHI antibody;Hyaluronate receptor antibody;IN antibody;INLU-related p80 Glycoprotein antibody;MC56 antibody;MDU2 antibody;MDU3 antibody;MGC10468 antibody;MIC4 antibody;MUTCH I antibody;MUTCH1 antibody;PGP-1 antibody;PGP-I antibody;PGP1 antibody;Phagocytic glycoprotein 1 antibody;Phagocytic glycoprotein I antibody;Soluble CD44 antibody;


Gene ID	960.0
SwissProt ID	P16070
Immunogen	A synthetic peptide of human CD44

Background

Cell localization:Membrane.The protein encoded by this gene is a cell-surface glycoprotein involved in cell-cell interactions, cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs). This protein participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing, hematopoiesis, and tumor metastasis. Transcripts for this gene undergo complex alternative splicing that results in many functionally distinct isoforms, however, the full length nature of some of these variants has not been determined. Alternative splicing is the basis for the structural and functional diversity of this protein, and may be related to tumor metastasis. [provided by RefSeq, Jul 2008],

Research Area

Image Data



Western blot analysis of lysates from A549 cells, using CD44 Rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG antibody was used to detect the antibody.