## **Product Name: CD38 Rabbit Monoclonal Antibody**

Catalog #: AMRe21443



#### **Summary**

Production Name CD38 Rabbit Monoclonal Antibody

**Description** Rabbit Monoclonal Antibody

**Host** Rabbit

**Application** WB,IHC,IF,IP,ELISA **Reactivity** Human,Mouse,Rat

#### **Performance**

ConjugationUnconjugatedModificationUnmodifiedIsotypeIgG,KappaClonalityMonoclonalFormLiquid

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

**Buffer** PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protective protein

**Purification** Protein A

### **Immunogen**

Gene Name CD38

CD38;ADP-ribosyl cyclase 1;Cyclic ADP-ribose hydrolase 1;CADPr hydrolase 1;T10;CD Alternative Names

antigen CD38

 Gene ID
 952.0

 SwissProt ID
 P28907.

### **Application**

IHC 1:200-1:1000;WB 1:1000-1:5000;IF 1:200-1:1000;ELISA 1:5000-1:20000;IP 1:50-

Dilution Ratio

1:200,

Molecular Weight Calculated MW:34kD;Observed MW:45kD

## **Background**


# Product Name: CD38 Rabbit Monoclonal Antibody Catalog #: AMRe21443



Cell localization:Membrane.The protein encoded by this gene is a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Crystal structure analysis demonstrates that the functional molecule is a dimer, with the central portion containing the catalytic site. It is used as a prognostic marker for patients with chronic lymphocytic leukemia. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015],

#### **Research Area**

#### **Image Data**



Western blot analysis of lysates from Mouse spleen

cells, using CD38 Rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG antibody was used to detect the antibody.

#### Note

For research use only.