Product Name: Chk2 Rabbit Monoclonal Antibody

Catalog #: AMRe21455



Summary

Production Name Chk2 Rabbit Monoclonal Antibody

Description Rabbit Monoclonal Antibody

Host Rabbit

Application WB,IHC,IF,IP,ELISA **Reactivity** Human,Mouse,Rat

Performance

ConjugationUnconjugatedModificationUnmodifiedIsotypeIgG,KappaClonalityMonoclonalFormLiquid

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protective protein

Purification Protein A

Immunogen

Gene Name CHEK2

CHEK2;CDS1;CHK2;RAD53;Serine/threonine-protein kinase Chk2;CHK2 checkpoint Alternative Names

homolog;Cds1 homolog;Hucds1;hCds1;Checkpoint kinase 2

 Gene ID
 11200.0

 SwissProt ID
 096017.

Application

IHC 1:200-1:1000;WB 1:1000-1:5000;IF 1:200-1:1000;ELISA 1:5000-1:20000;IP 1:50-

Dilution Ratio

1:200,

Molecular Weight Calculated MW:61kD;Observed MW:61kD

Background


Product Name: Chk2 Rabbit Monoclonal Antibody Catalog #: AMRe21455



Cell localization:Nucleus.In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutati

Research Area

Image Data



Western blot analysis of lysates from K562

cells, using Chk2 Rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG antibody was used to detect the antibody.

Note

For research use only.