

---

**Product Name: Iba1 Rabbit Monoclonal Antibody****Catalog #: AMRe21499**

For research use only.

**Summary**

<b>Description</b>	Recombinant rabbit monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC,ICC/IF,ELISA,IP
<b>Reactivity</b>	Human,Mouse,Rat
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG,Kappa
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Concentration</b>	0.3mg/ml. The concentration of this product may be batch-dependent.
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%protective protein
<b>Purification</b>	Protein A

**Application**

<b>Dilution Ratio</b>	WB 1:1000-1:5000,IHC 1:2000-1:4000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200
<b>Molecular Weight</b>	Calculated MW:17kD;Observed MW:17kD

**Antigen Information**

<b>Gene Name</b>	AIF AIF 1;AIF-1;Aif1;AIF1 protein;AIF1_HUMAN;Allograft inflammatory factor 1;Allograft inflammatory factor 1 splice variant G;allograft inflammatory factor-1 splice variant Hara-1;balloon angioplasty responsive transcription;BART 1;G1;G1 putative splice variant of
<b>Alternative Names</b>	allograft inflammatory factor 1;IBA 1;IBA1;interferon gamma responsive transcript;Interferon responsive transcript 1;interferon responsive transcript factor 1;ionized calcium binding adapter molecule 1;ionized calcium-binding adapter molecule 1;ionized calcium-binding adapter molecule;IRT 1;IRT1;Microglia response factor;MRF1;Protein g1;
<b>Gene ID</b>	199.0


<b>SwissProt ID</b>	P55008
<b>Immunogen</b>	A synthetic peptide of human Iba1

## Background

Cell localization: Cytoplasm. This gene encodes a protein that binds actin and calcium. This gene is induced by cytokines and interferon and may promote macrophage activation and growth of vascular smooth muscle cells and T-lymphocytes. Polymorphisms in this gene may be associated with systemic sclerosis. Alternative splicing results in multiple transcript variants, but the full-length and coding nature of some of these variants is not certain. [provided by RefSeq, Jan 2016],

## Research Area


## Image Data



Western blot analysis of lysates from Mouse spleen cells, using Iba1 Rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG antibody was used to detect the antibody.



Immunohistochemical analysis of paraffin-embedded Human liver tissue. 1, Iba1 Rabbit Monoclonal Antibody was diluted at 1:200(4°C, overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C, 20min). 3, Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Mouse liver tissue. 1, Iba1 Rabbit Monoclonal Antibody was diluted at 1:200(4°C, overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C, 20min). 3, Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Rat liver tissue. 1, Iba1 Rabbit Monoclonal Antibody was diluted at 1:200(4°C, overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C, 20min). 3, Secondary antibody was diluted at 1:200(room temperature, 30min).

