

**Product Name: TATA Box Binding Protein Rabbit
Monoclonal Antibody
Catalog #: AMRe21552**

Summary

Production Name	TATA Box Binding Protein Rabbit Monoclonal Antibody
Description	Recombinant rabbit monoclonal antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA,IP
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG,Kappa
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%protective protein
Purification	Protein A

Immunogen

Gene Name	TBP TBP;GTF2D1;TF2D;TFIID;TATA-box-binding protein;TATA sequence-binding
Alternative Names	protein;TATA-binding factor;TATA-box factor;Transcription initiation factor TFIID TBP subunit
Gene ID	6908.0
SwissProt ID	P20226.

Application

Dilution Ratio	WB 1:1000-1:5000,IHC 1:400-1:1000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200
-----------------------	---

**Product Name: TATA Box Binding Protein Rabbit
Monoclonal Antibody
Catalog #: AMRe21552**

Molecular Weight


Calculated MW:38kD;Observed MW:38kD

Background

Cell localization:Nucleus.Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes TBP, the TATA-binding protein. A distinctive feature of TBP is a long string of glutamines in the N-terminus. This region of the protein modulates the DNA bin

Research Area


Image Data




Western blot analysis of lysates from HEK293

cells, using TATA Box Binding Protein Rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG antibody was used to detect the antibody.


**Product Name: TATA Box Binding Protein Rabbit
Monoclonal Antibody
Catalog #: AMRe21552**



Immunohistochemical analysis of paraffin-embedded human colon carcinoma tissue. 1, primary Antibody was diluted at 1:200(4°C,overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded mouse colon tissue. 1, primary Antibody was diluted at 1:200(4°C,overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded rat colon tissue. 1, primary Antibody was diluted at 1:200(4°C,overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

**Product Name: TATA Box Binding Protein Rabbit
Monoclonal Antibody
Catalog #: AMRe21552**



Note

For research use only.