

**Product Name: FTCD Recombinant Human Monoclonal Antibody**  
**Catalog #: AMRe83675**

---



## Summary

<b>Production Name</b>	FTCD Recombinant Human Monoclonal Antibody
<b>Description</b>	Recombinant antibody
<b>Host</b>	Human Cell
<b>Application</b>	WB,IHC
<b>Reactivity</b>	Human

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	Human IgG1
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Purified antibody in PBS with 0.05% sodium azide
<b>Purification</b>	Affinity Purification

## Immunogen

<b>Gene Name</b>	FTCD
<b>Alternative Names</b>	FTCD
<b>Gene ID</b>	
<b>SwissProt ID</b>	O95954.Purified recombinant fragment of human FTCD

## Application

<b>Dilution Ratio</b>	WB:1:500-1:1000,IHC:1:100-1:200
<b>Molecular Weight</b>	58.9kDa

## Background


---

**Product Name: FTCD Recombinant Human Monoclonal Antibody**  
**Catalog #: AMRe83675**




## Research Area

## Image Data




96-well microtitre plates were coated with 1 µg/mL of FTCD protein at 4 °C overnight. Plates were blocked with 2 % BSA in PBS-T (0.1 % Tween) at room temperature for 1 hour. Serial diluted of AMRe83675 was used as the primary antibody and added into the plates for 1 hour incubation. Plates were washed with PBS-T for three times and then incubated with HRP conjugated Human IgG1 for 1 hour. After washed with PBS-T for three times, the plates were incubated with ABTS for 15 minute. The absorbance (optical density) was read with a plate reader.



Western blot analysis using FTCD mAb against Hep G2 lysate

**Product Name: FTCD Recombinant Human Monoclonal Antibody**  
**Catalog #: AMRe83675**

---



IHC-P analysis of liver tissue by FTCD antibody (AMRe83675)

### **Note**

For research use only.