

### **Product Name: TRAF6BP Rabbit Monoclonal Antibody**

Catalog #: AMRe84207

For research use only.

#### **Summary**

**Description** Recombinant rabbit monoclonal antibody

Host Rabbit
Application WB,ICC

Reactivity Human,Mouse,Rat
Conjugation Unconjugated
Modification Unmodified

**Isotype** IgG

Clonality Monoclonal
Form Liquid

Concentration 0.39mg/ml. The concentration of this product may be batch-dependent.

Storage Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

Shipping Ice bags

Purified antibody in PBS with 0.05% sodium azide,0.05% protective protein and 50% **Buffer** 

glycerol.

**Purification** Affinity Purification

### **Application**

**Dilution Ratio** WB 1:1000-1:2000,ICC 1:50-1:200

Molecular Weight 91 kDa

# **Antigen Information**

Gene Name TRAF6BP

CALCOCO3; D6Ertd404e; D6Ertd772e; PRO0105; T6BP; TAX1BP1; tax1bp1b; TXBP151;;Tax1

Alternative Names binding protein 1

Gene ID

SwissProt ID Q86VP1

**Immunogen** A synthesized peptide derived from human Tax1 binding protein 1

## **Background**

Ubiquitin-binding adapter that participates in inflammatory, antiviral and innate immune processes as well as selective


Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




autophagy regulation. Plays a key role in the negative regulation of NF-kappa-B and IRF3 signalings by acting as an adapter for the ubiquitin-editing enzyme A20/TNFAIP3 to bind and inactivate its substrates.

#### **Research Area**


# **Image Data**



Western blot analysis of TRAF6BP expression in HepG2 cell lysate.




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.