

---

**Product Name: HLTF Rabbit Monoclonal Antibody****Catalog #: AMRe84247**

For research use only.

**Summary**

<b>Description</b>	Recombinant rabbit monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC,ICC/IF,ICC
<b>Reactivity</b>	Human,Rat
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Concentration</b>	0.71mg/ml. The concentration of this product may be batch-dependent.
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Purified antibody in PBS with 0.05% sodium azide,0.05% protective protein and 50% glycerol.
<b>Purification</b>	Affinity Purification

**Application**

<b>Dilution Ratio</b>	WB 1:1000-1:2000,IHC 1:100-1:200,ICC/IF 1:50-1:200,ICC 1:50-1:200
<b>Molecular Weight</b>	Calculated MW: 114 kDa ; Observed MW: 120 kDa

**Antigen Information**

<b>Gene Name</b>	HLTF
<b>Alternative Names</b>	HIP116; HIP116A; HLTF 1; Hltf; HLTF1; p113; RNF80; SMARC A3; SMARCA 3; SMARCA3; SNF2L3; ZBU1;;HLTF
<b>Gene ID</b>	
<b>SwissProt ID</b>	Q14527
<b>Immunogen</b>	A synthesized peptide derived from human HLTF


**Background**

Has both helicase and E3 ubiquitin ligase activities. Possesses intrinsic ATP-dependent nucleosome-remodeling activity; This

activity may be required for transcriptional activation or repression of specific target promoters (By similarity).

## Research Area

## Image Data



Western blot analysis of HLTF expression in K562 cell lysate.

All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.

