

---

**Product Name: Gli3 Rabbit Monoclonal Antibody****Catalog #: AMRe84339**

For research use only.

**Summary**

|                      |   |
|----------------------|---|
| <b>Description</b>   | Recombinant rabbit monoclonal antibody  |
| <b>Host</b>          | Rabbit  |
| <b>Application</b>   | WB,ICC  |
| <b>Reactivity</b>    | Human   |
| <b>Conjugation</b>   | Unconjugated  |
| <b>Modification</b>  | Unmodified  |
| <b>Isotype</b>       | IgG   |
| <b>Clonality</b>     | Monoclonal  |
| <b>Form</b>          | Liquid  |
| <b>Concentration</b> | 0.71mg/ml. The concentration of this product may be batch-dependent.                        |
| <b>Storage</b>       | Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.                 |
| <b>Shipping</b>      | Ice bags  |
| <b>Buffer</b>        | Purified antibody in PBS with 0.05% sodium azide,0.05% protective protein and 50% glycerol. |
| <b>Purification</b>  | Affinity Purification   |

**Application**

|                         |   |
|-------------------------|---|
| <b>Dilution Ratio</b>   | WB 1:1000-1:2000,ICC 1:50-1:200               |
| <b>Molecular Weight</b> | Calculated MW: 170 kDa ; Observed MW: 190 kDa |

**Antigen Information**

|                          |   |
|--------------------------|---|
| <b>Gene Name</b>         | Gli3  |
| <b>Alternative Names</b> | ACLS; GCPS; GLI3; GLI3FL; PAPA; PAPA1; PAPB; PHS; PPDIV;;GLI3 |
| <b>Gene ID</b>           |   |
| <b>SwissProt ID</b>      | P10071  |
| <b>Immunogen</b>         | A synthesized peptide derived from human GLI3                 |


**Background**

Has a dual function as a transcriptional activator and a repressor of the sonic hedgehog (Shh) pathway, and plays a role in limb development. The full-length GLI3 form (GLI3FL) after phosphorylation and nuclear translocation, acts as an activator (GLI3A)


while GLI3R, its C-terminally truncated form, acts as a repressor.

## Research Area

## Image Data



Western blot analysis of Gli3 expression in 293 cell lysate.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.