

**Product Name: Ube2N/Ubc13 Rabbit Monoclonal Antibody****Catalog #: AMRe84356**

For research use only.

**Summary**

<b>Description</b>	Recombinant rabbit monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC,ICC/IF,ICC
<b>Reactivity</b>	Human,Mouse,Rat
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Concentration</b>	
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Purified antibody in PBS with 0.05% sodium azide,0.05% protective protein and 50% glycerol.
<b>Purification</b>	Affinity Purification

**Application**

<b>Dilution Ratio</b>	WB 1:1000-1:2000,IHC 1:100-1:200,ICC/IF 1:50-1:200,ICC 1:50-1:200
<b>Molecular Weight</b>	17 kDa

**Antigen Information**

<b>Gene Name</b>	Ube2N/Ubc13
<b>Alternative Names</b>	BLU; HEL-S-71; Ubc13; UbCH ben; UbCH13; UBCHBEN; Ube2n;;UBE2N
<b>Gene ID</b>	
<b>SwissProt ID</b>	P61088
<b>Immunogen</b>	A synthesized peptide derived from human UBE2N


**Background**

The UBE2V1-UBE2N and UBE2V2-UBE2N heterodimers catalyze the synthesis of non-canonical 'Lys-63'-linked polyubiquitin chains. This type of polyubiquitination does not lead to protein degradation by the proteasome. Mediates transcriptional

activation of target genes. Plays a role in the control of progress through the cell cycle and differentiation.

## Research Area

## Image Data



Western blot analysis of Ube2N / Ubc13 expression in Daudi cell lysate.