

---

**Product Name: Histone H3(mutatedK27Met) Rabbit Monoclonal Antibody****Catalog #: AMRe84538**

For research use only.

**Summary**

<b>Description</b>	Recombinant rabbit monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,ICC,IP
<b>Reactivity</b>	Human
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Concentration</b>	
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Purified antibody in PBS with 0.05% sodium azide,0.05% protective protein and 50% glycerol.
<b>Purification</b>	Affinity Purification

**Application**

<b>Dilution Ratio</b>	WB 1:1000-1:2000,ICC 1:50-1:200,IP 1:20-1:50
<b>Molecular Weight</b>	Calculated MW: 15 kDa ; Observed MW: 18 kDa

**Antigen Information**

<b>Gene Name</b>	Histone H3(mutatedK27Met)
<b>Alternative Names</b>	Histone H3.1, Histone H3, HIST1H3A;;Histone H3 (K27M mutated)
<b>Gene ID</b>	
<b>SwissProt ID</b>	P68431
<b>Immunogen</b>	A synthesized peptide derived from human Histone H3.1 (K27M mutated)


**Background**

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA

repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

## Research Area

## Image Data



Western blot analysis of Histone H3 (mutated K27 Met) expression in recombinant protein lysate.