

Product Name: Histone H2B(formylK108) Rabbit Monoclonal Antibody Catalog #: AMRe84632

For research use only.

Summary

Description Recombinant rabbit monoclonal antibody

Host Rabbit
Application WB,IHC

Reactivity Human,Mouse,Rat
Conjugation Unconjugated
Modification Unmodified

Isotype IgG

Clonality Monoclonal
Form Liquid

Concentration 0.62mg/ml. The concentration of this product may be batch-dependent.

Storage Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

Shipping Ice bags

Purified antibody in PBS with 0.05% sodium azide,0.05% protective protein and 50% **Buffer**

glycerol.

Purification Affinity Purification

Application

Dilution Ratio WB 1:1000-1:2000,IHC 1:100-1:200

Molecular Weight 14 kDa

Antigen Information

Gene Name Histone H2B(formylK108)

Alternative Names Histone H2B;;Formyl-Histone H2B type 2E (K109)

Gene ID

SwissProt ID Q16778

A synthesized peptide derived from human Histone H2B type 2E around the formylation site **Immunogen**

of K109

Background

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the


Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Research Area


Image Data



Western blot analysis of Histone H2B (formyl K108) expression in Hela cell lysate.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838