

---

**Product Name: HTF9C Rabbit Monoclonal Antibody****Catalog #: AMRe84638**

For research use only.

**Summary**

<b>Description</b>	Recombinant rabbit monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC,ICC/IF,ICC
<b>Reactivity</b>	Human
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Concentration</b>	0.71mg/ml. The concentration of this product may be batch-dependent.
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Purified antibody in PBS with 0.05% sodium azide,0.05% protective protein and 50% glycerol.
<b>Purification</b>	Affinity Purification

**Application**

<b>Dilution Ratio</b>	WB 1:1000-1:2000,IHC 1:100-1:200,ICC/IF 1:50-1:200,ICC 1:50-1:200
<b>Molecular Weight</b>	Calculated MW: 69 kDa ; Observed MW: 75 kDa

**Antigen Information**


<b>Gene Name</b>	HTF9C
<b>Alternative Names</b>	HTF9C; TRMT2A;;TRMT2A
<b>Gene ID</b>	
<b>SwissProt ID</b>	Q8IZ69
<b>Immunogen</b>	A synthesized peptide derived from human TRMT2A

**Background**


S-adenosyl-L-methionine-dependent methyltransferase that catalyzes the formation of 5-methyl-uridine in tRNAs and some mRNAs. Mainly catalyzes the methylation of uridine at position 54 (m5U54) in cytosolic tRNAs.

## Research Area

## Image Data



Western blot analysis of HTF9C expression in A431 cell lysate.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.