

---

**Product Name: FANCD2 Rabbit Monoclonal Antibody****Catalog #: AMRe86270**

For research use only.

**Summary**

<b>Description</b>	Recombinant rabbit monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC,ICC/IF,FC,IP
<b>Reactivity</b>	Human,Mouse,Rat
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Concentration</b>	0.55mg/ml. The concentration of this product may be batch-dependent.
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% protective protein. Stable for 12 months from date of receipt.
<b>Purification</b>	Affinity Purification

**Application**

<b>Dilution Ratio</b>	WB 1:1000-1:5000,IHC 1:50-1:200,ICC/IF 1:200-1:500,FC 1:100-1:200,IP 1:10-1:100
<b>Molecular Weight</b>	Calculated MW:164 kDa; Observed MW:164 kDa

**Antigen Information**

<b>Gene Name</b>	FANCD2
<b>Alternative Names</b>	FA4; FAD; FACD; FAD2; FA-D2; FANCD
<b>Gene ID</b>	2177
<b>SwissProt ID</b>	Q9BXW9
<b>Immunogen</b>	A synthetic peptide of human FANCD2


**Background**

The Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCI (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The


previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group D2. This protein is monoubiquitinated in response to DNA damage, resulting in its localization to nuclear foci with other proteins (BRCA1 AND BRCA2) involved in homology-directed DNA repair. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2016]

## Research Area


### Image Data



Western blot analysis of extracts from HeLa cells using FANCD2 Rabbit Monoclonal Antibody at 1:1000.



Western blot analysis of extracts from HeLa, U2OS, A2780 cells using AMRe86270 at 1:500.



Immunofluorescence analysis of HeLa cells labelling FANCD2 with AMRe86270.