Product Name: NFkB p105/p50 Rabbit Monoclonal

Antibody

Catalog #: AMRe87472



Summary

Production Name NFkB p105/p50 Rabbit Monoclonal Antibody

Description Rabbit Monoclonal antibody

Host Rabbit

Application WB, IHC-P

Reactivity Human, Mouse, Rat

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

Clonality Monoclonal Form Liquid

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide

and 0.05% protective protein. Stable for 12 months from date of receipt.

Purification Affinity Purification

Immunogen

Buffer

Gene Name NFkB p105/p50

p50; KBF1; p105; EBP-1; CVID12; NF-kB1; NFKB-p50; NFkappaB; NF-kappaB; NFKB-

Alternative Names p105; NF-kappa-B

Gene ID 4790, 18033, 81736

SwissProt ID P19838, P25799, Q63369.

Application

Dilution Ratio WB: 1:1000-1:5000 IHC-P: 1:200-1:500

Molecular Weight Calculated MW:105 kDa; Observed MW:105,50 kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: NFkB p105/p50 Rabbit Monoclonal

Antibody

Catalog #: AMRe87472




Background

This gene encodes a 105 kD protein which can undergo cotranslational processing by the 26S proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFKB) protein complex. NFKB is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NFKB translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NFKB has been associated with a number of inflammatory diseases while persistent inhibition of NFKB leads to inappropriate immune cell development or delayed cell growth. Alternative splicing results in multiple transcript variants encoding different isoforms, at least one of which is proteolytically processed. [provided by RefSeq, Feb 2016]

Research Area

Image Data



Western blot analysis of extracts from PC-12 cells using AMRe87472 at 1:1000.

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838